

9. *Pogonia ophioglossoides*, Nutt. Near Boy pond and Elbow pond, not common.

The *Habenarias* are well represented in this region and I meet most of the following species every season while tramping along the brooks.

10. *Habenaria tridentata*, Hook. Below the railroad bridge, abundant; frequent in the Pondfield meadows.

11. *Habenaria hyperborea*, R. Br. Likely to be met with anywhere in wet places on the slopes of the mountains forming the valley. At Beaver Falls, on the eastern slope of Moosilauke, I found hundreds of this stately plant in a wild, deep ravine on August 1, 1895, and I named one portion of the falls, "Orchid Falls."

12. *Habenaria dilata*, Gray. Frequent near Echo Lake and the Basin in the Notch.

13. *Habenaria Hookeri*, Torr. On the Warren road, amidst dense mats of *Linnaea borealis*. I have found it once in August, 1896, and once this year — only one specimen badly worm-eaten.

14. *Habenaria orbiculata*, Torr. Very common in old logging roads in the woods and deep forests; from two to a dozen are frequently found near each other.

15. *Habenaria fimbriata*, R. Br. This beautiful and showy orchid is common in the low grass lands along the river near thickets.

16. *Cypripedium pubescens*, Willd. Frequent in cold and shady bogs among the mountains.

17. *Cypripedium acaule*, Ait. Very abundant with *Epigaea repens* on the hillsides and ledges bordering the Pemigewasset.

---

HYDRASTIS CANADENSIS, L. IN VERMONT. — Last June Mr. F. H. Horsford of Charlotte informed me that one of his collectors of bulbs had brought in specimens of *Hydrastis Canadensis*. On August 8, after Mr. Horsford had kindly given me the necessary directions, I visited the station, which is in the town of Shelburne, and found a colony of twenty-five or thirty plants, from which I ventured to take three specimens. This is apropos of Mr. Bissell's note in RHODORA, vol. I, p. 157, regarding the occurrence of this species in Connecticut. — EZRA BRAINERD, Middlebury, Vermont.